

PUBLICATION LISTS OF MASAHIRO NAKAOKA

Papers published in peer-reviewed journals

- 1: Nakaoka, M. (1992a) Age determination and growth analysis based on external shell rings of the protobranch bivalve *Yoldia notabilis* Yokoyama in Otsuchi Bay, northeastern Japan. *Benthos Research* **43**: 53-66
- 2: Nakaoka, M. (1992b) Spatial and seasonal variation in growth rate and secondary production of *Yoldia notabilis* in Otsuchi Bay, Japan, with reference to the influence of food supply from the water column. *Marine Ecology Progress Series* **88**: 215-223
- 3: Nakaoka, M. (1993) Yearly variation in recruitment and its effect on population dynamics in *Yoldia notabilis* (Mollusca: Bivalvia), analyzed using projection matrix model. *Researches on Population Ecology* **35**: 199-213
- 4: Nakaoka, M. (1994) Size-dependent reproductive traits of *Yoldia notabilis* (Bivalvia: Protobranchia). *Marine Ecology Progress Series* **114**: 129-137
- 5: Nakaoka, M. and Matsui S. (1994) Annual variation in the growth rate of *Yoldia notabilis* (Bivalvia: Nuculanidae) in Otsuchi Bay, northeastern Japan, analyzed using shell microgrowth patterns. *Marine Biology* **119**: 397-404
- 6: Nakaoka, M. (1996a) Size-dependent survivorship of the bivalve *Yoldia notabilis* (Yokoyama, 1920): the effect of crab predation. *Journal of Shellfish Research* **15**: 355-362
- 7: Nakaoka, M. (1996b) Dynamics of age- and size-structured populations in fluctuating environments: Applications of stochastic matrix models to natural populations. *Researches on Population Ecology* **38**: 141-152
- 8: Nakaoka, M. (1997) Demography of the marine bivalve *Yoldia notabilis* in fluctuating environments: an analysis using stochastic matrix model. *Oikos* **79**: 59-68
- 9: Nakaoka, M. (1998) Optimal resource allocation of the marine bivalve *Yoldia notabilis*: the effects of size-limited reproductive capacity and size-dependent mortality. *Evolutionary Ecology* **12**: 347-361
- 10: Nakaoka, M. and Ohta, S. (1998) Seasonal variation in reproduction and population structure of the bivalve *Limopsis tajimae* in the upper bathyal zone of Suruga Bay, Japan. *Marine Biology* **132**: 471-481
- 11: Komatsu, T., Nakaoka, M. and Kawai, H. (1999) Development of evaluation method of impacts by heavy-oil pollution on inter-tidal ecosystem using Geographical Information System (GIS). *Bulletin on Coastal Oceanography* **37**: 35-39
- 12: Nakaoka, M. and Aioi, K. (1999) Growth of the seagrass *Halophila ovalis* at the dugong trails compared to existing within-patch variation in a Thailand intertidal flat. *Marine Ecology Progress Series* **184**: 97-103
- 13: Toyohara, T., Nakaoka, M. and Aioi, K. (1999) Population dynamics and reproductive traits of phytal gastropods in seagrass bed in Otsuchi Bay, northeastern Japan. *P. S. Z. N. Marine Ecology* **20**: 273-290
- 14: Nakaoka, M. (2000) Nonlethal effects of predators on prey populations: predator-mediated change in bivalve growth. *Ecology* **81**: 1031-1045
- 15: Aioi, K., Nakaoka, M., Kouchi, N. and Omori, Y. (2000) A new record of *Zostera asiatica* Miki (Zosteraceae) in Funakoshi Bay, Iwate Prefecture. *Otsuchi Marine Science* **25**: 23-26
- 16: Nakaoka, M., Kouchi, N. and Aioi, K. (2000) Growth and shoot dynamics of *Zostera caulescens* Miki in Funakoshi Bay, Japan: how does it maintain high canopy structure? *Biologia Marina Mediterranea* **7**: 103-106
- 17: Watanabe, M., Nakaoka, M. and Mukai, H. (2000) Growth and productivity of *Zostera asiatica* and *Zostera marina* in Akkeshi, northern Japan. *Biologia Marina Mediterranea* **7**: 156-159
- 18: Kouchi, N. and Nakaoka, M. (2000) Distribution of encrusting bryozoa on *Zostera caulescens* in Funakoshi Bay, Japan: effects of seagrass vertical structure on epifauna. *Biologia Marina Mediterranea* **7**: 247-250
- 19: Komatsu, T. and Nakaoka, M. (2000) Water flow and sedimentation rate in seagrass beds off Khao Baena in Trang Province, Thailand. *Biologia Marina Mediterranea* **7**: 240-242
- 20: Mukai, H., Aioi, K., Lewmanomont, K., Matsumasa, M., Nakaoka, M., Nojima, S. Supanwanid,

- C., Suzuki, T., and Toyohara, T. (2000) Dugong grazing on *Halophila* beds in Haad Chao Mai National Park. *Biologia Marina Mediterranea* **7**: 268-270
- 21: Nakaoka, M. and Iizumi, H. (2000) Magnitude of within-patch variation in seagrass *Halophila ovalis* growth affected by adjacent *Thalassia hemprichii* vegetation. *Ecological Research* **15**: 415-424
- 22: Nakaoka, M. and Toyohara, T. (2000) Effects of seagrass patch structure on the mobile epifaunal community in a subtidal seagrass meadow in Thailand. *Benthos Research* **55**: 53-61
- 23: Nakaoka, M. and Aioi, K. (2001) Ecology of seagrasses *Zostera* spp. (Zosteraceae) in Japanese waters: A review. *Otsuchi Marine Science* **26**: 7-22
- 24: Toyohara, T., Nakaoka, M. and Tsuchida, E. (2001) Population dynamics and life history traits of *Siphonacmea oblongata* Yokohama on seagrass leaf in Otsuchi Bay (Siphonariidae, Pulamonata). *Venus (Japanese Journal of Malacology)* **60** (1/2): 27-36
- 25: Nakaoka, M. (2001) Small-scale variation in a benthic community at an intertidal flat in Thailand: effects of spatial heterogeneity of seagrass vegetation. *Benthos Research* **56**: 63-71
- 26: Nakaoka, M., Toyohara, T. and Matsumasa, M. (2001) Seasonal and between-substrate variation in mobile epifaunal community in a multispecific seagrass bed of Otsuchi Bay, Japan. *P. S. Z. N. Marine Ecology* **22**: 379-395
- 27: Nakaoka, M. (2002) Predation on seeds of seagrasses *Zostera marina* and *Zostera caulescens* by a tanaid crustacean *Zeuxo* sp. *Aquatic Botany* **72**: 99-106
- 28: Tanaka, N., Omori, Y., Nakaoka, M. and Aioi, K. (2002) Gene flow among populations of *Zostera caespitosa* Miki (Zosteraceae) in Sanriku Coast, Japan. *Otsuchi Marine Science* **27**: 17-22
- 29: Nakaoka, M., Mukai, H. and Chunhabundit, S. (2002) Impacts of dugong foraging on benthic animal communities in a Thailand seagrass bed. *Ecological Research* **17**: 625-638
- 30: Komatsu, T., Igarashi, C., Tatsukawa, K., Nakaoka, M. and Taira, A. (2002) Mapping of seagrass and seaweed beds using hydro-acoustic methods. *Fisheries Science* **68** sup. I: 580-583
- 31: Komatsu, T., Nakaoka, M. and Sugimoto, T. (2003) Changes in coastal ecosystem due to modifications of land use pattern. *Bulletin on Coastal Oceanography* **40**: 149-157
- 32: Nakaoka, M., Aioi, K., Kouchi, N., Omori, Y., Tanaka, N. and Tatsukawa, K. (2003) Distribution, productivity, life history and biodiversity of seagrass community along Sanriku Coast: A review. *Otsuchi Marine Science* **28**: 31-38
- 33: Komatsu T., Nakaoka, M., Kawai H., Yamamoto, T., Marine Life Research Group of Takeno and Ohwada, K. (2003) Impacts of the Nakhodka heavy-oil spill on an intertidal ecosystem: An approach to impact evaluation using GIS. *Marine Pollution Bulletin* **47**: 99-104
- 34: Yamamoto, T., Nakaoka, M., Komatsu T., Kawai H., Marine Life Research Group of Takeno and Ohwada, K. (2003) Impacts by heavy-oil spill from the Russian tanker *Nakhodka* on intertidal ecosystems: Recovery of animal community. *Marine Pollution Bulletin* **47**: 91-98
- 35: Tanaka, N., Kuo, J., Omori, Y., Nakaoka, M. and Aioi, K. (2003) Phylogenetic relationships in the genera *Zostera* and *Heterozostera* (Zosteraceae) based on matK sequence data. *Journal of Plant Research* **116**: 273-279
- 36: Nakaoka, M., Kouchi, N. and Aioi, K. (2003) Seasonal dynamics of *Zostera caulescens*: relative importance of flowering shoots to net production. *Aquatic Botany* **77**: 277-293
- 37: Yoshida, M., Kouchi, N. and Nakaoka, M. (2003) Citizens' Involvement for Seagrass Watch Survey in Okinawa Island. *Japanese Journal of Conservation Ecology* **8**: 119-128
- 38: Komatsu, T., Umezawa, Y., Nakaoka, M., Supanwanid, C. and Kanamoto, Z. (2004) Water flow and sediment in *Enhalus acoroides* and other seagrass beds in the Andaman Sea, off Khao Bae Na, Thailand. *Coastal Marine Science* **29**: 63-68
- 39: Nakaoka, M., Tanaka, Y. and Watanabe M. (2004) Species diversity and abundance of seagrasses in southwestern Thailand under different influence of river discharge. *Coastal Marine Science* **29**: 75-80
- 40: Okuda, T., Noda, T., Yamamoto, T. Ito, N. and Nakaoka, M. (2004) Latitudinal gradient of species diversity: multi-scale variability in rocky intertidal sessile assemblages along the Northwestern Pacific coast. *Population Ecology* **46**: 159-170

- 41: Tanaka, Y. and Nakaoka, M. (2004) Emergence stresses and morphological constraints affect the species distribution and growth of subtropical intertidal seagrasses. *Marine Ecology Progress Series* **284**: 117-131
- 42: Noda, T., Fukatsu, Y., Okuda, T., Hori, M., Nakaoka, M. and Yamamoto, T. (2005) Seven forms of rarity in rocky intertidal algal assemblages along the Northwestern Pacific coast of Japan. *Japanese Journal of Conservation Ecology* **10**: 1-10 (in Japanese with English abstract)
- 43: Nakaoka, M. (2005) Plant-animal interactions in seagrass beds: ongoing and future challenges for understanding population and community dynamics. *Population Ecology* **47**: 167-177
- 44: Watanabe, M., Nakaoka, M. Mukai, H. (2005) Seasonal variation in vegetative growth and production of the endemic seagrass *Zostera asiatica* in Japan: a comparison with sympatric *Zostera marina*. *Botanica Marina* **48**: 266-273
- 45: Yamakita T., Nakaoka M., Kondoh A, Ishii M, Shoji Y (2005) Long-term spatial dynamics of a seagrass bed on Futtsu tidal flat in Tokyo Bay. *Japanese Journal of Conservation Ecology* (in Japanese with English abstract) **10**: 129-138
- 46: Tanaka, N., Shoji, Y., Nakaoka, M., Ishii, M., Lim, B. K. (2006) Distribution of *Zostera marina* (Zosteraceae) in coastal seawaters in the Sagami Sea. *Memoirs of the National Science Museum* **42**: 53-57
- 47: Nakaoka M., Ito N, Yamamoto T, Okuda T, Noda T (2006) Similarity of rocky intertidal assemblages along the Pacific Coast of Japan: Effects of spatial scales and geographic distance. *Ecological Research* **21**: 425-435
- 48: Kouchi, N., Nakaoka M., Mukai, H. (2006) Effects of temporal dynamics and vertical structure of the seagrass *Zostera caulescens* on distribution and recruitment of the epifaunal encrusting bryozoa *Microporella trigonellata*. *Marine Ecology* **27**: 145-153
- 49: Tanaka, Y., Nakaoka, M. (2006) Morphological variation of tropical seagrasses, *Cymodocea serrulata* and *C. rotundata* in response to sediment condition and light attenuation. *Botanica Marina* **49**: 365-371
- 50: Kawai H., Kamiya, M., Komatsu T., Nakaoka, M., Yamamoto, T. and the Marine Life Research Group of Takeno (2007) Ten years' monitoring of intertidal macroalgal vegetation of Hyogo Prefecture, northwestern coast of Honshu, Japan to assess the impact of the *Nakhodka* oil spill. *Algae* **22**: 1-10
- 51: Yoshida, M., Kouchi, N. and Nakaoka, M. (2007) Spatial dynamics of seagrass beds analyzed by integrated approach using remote sensing and citizen-based monitoring. *Japanese Journal of Conservation Ecology* **12**: 10-19
- 52: Hori, M., Kamimura, S. and Nakaoka, M. (2007) The Importance of Ecosystem-Functioning Research for Conservation and Sustainable Utilization of Coastal Seas. *Japanese Journal of Benthology* **62**: 68-72
- 53: Nakaoka, M., Watanabe, K., Era, T. and Ishii, M. (2007) Evaluation of relationships between biodiversity and ecosystem functions in coastal seas: a case study of seagrass beds in Tokyo Bay. *Japanese Journal of Benthology* **62**: 82-87
- 54: Kamimura, S., Hori, M. and Nakaoka, M. (2007) An approach to conservation and sustainable use in coastal ecosystems: confronting the real issues and the solutions. *Japanese Journal of Benthology* **62**: 98-103
- 55: Yamada, K., Hori, M., Tanaka, Y., Hasegawa, Y. and Nakaoka, M. (2007) Temporal and spatial macrofaunal community changes along a salinity gradient in seagrass meadows of Akkeshi-ko estuary and Akkeshi Bay, northern Japan. *Hydrobiologia* **592**: 345-358
- 56: Nakaoka, M., Tanaka, Y., Mukai, H., Suzuki, T. and Aryuthaka C. (2007) Tsunami impacts on biodiversity of seagrass communities in the Andaman Sea, Thailand: (1) Seagrass abundance and diversity. In, Rigby P. R., Shirayama, Y., editors, Selected Papers of the NaGISA World Congress 2006, pp. 49-56
- 57: Whanpetch, N., Nakaoka, M., Mukai, H., Suzuki, T., Nojima, S., Kawai, T., and Aryuthaka, C. (2007) Tsunami impacts on biodiversity of seagrass communities in the Andaman Sea, Thailand: (2) Abundance and diversity of benthic animals. In, Rigby P. R., Shirayama, Y., editors, Selected Papers of the NaGISA World Congress 2006, pp. 57-66
- 58: Tanaka, Y. and Nakaoka, M. (2007) Interspecific variation in photosynthesis and respiration

- balance of three seagrasses in relation to light availability. *Marine Ecology Progress Series* **350**:63-70
- 59: Nakaoka, M., Matsumasa, M., Toyohara, T. and Williams, S.L. (2008) Animals on marine flowers: Does the presence of flowering shoots affect mobile epifaunal assemblage in an eelgrass meadow? *Marine Biology* **153**: 589-598
- 60: Okuda, T., Noda, T., Yamamoto, T., Hori, M. and Nakaoka, M. (2009) Latitudinal gradients in species richness in assemblages of sessile animals in rocky intertidal zone: mechanisms determining scale-dependent variability. *Journal of Animal Ecology* **78**: 328-337
- 61: Yamakita, T. and Nakaoka, M. (2009) Scale dependency in seagrass dynamics: how does the neighboring effect vary with grain of observation? *Population Ecology* **51**: 33-40
- 62: Hori, M., Suzuki, T., Monthum, Y., Srisombat, T., Tanaka, Y., Nakaoka, M. and Mukai, H. (2009) High seagrass diversity and canopy height increase the associated fish diversity and abundance. *Marine Biology* **156**: 1447-1458
- 63: Tsujino, M., Hori, M., Okuda, T., Nakaoka, M., Yamamoto, T. and Noda, T. (2010) Distance decay of community dynamics in rocky intertidal sessile assemblages evaluated by transition matrix models. *Population Ecology* **52**: 171-180
- 64: Yamada, K., Hori, M., Tanaka, Y., Hasegawa, N. and Nakaoka, M. (2010) Contribution of different functional groups to the diet of major predatory fishes at a seagrass meadow in northeastern Japan. *Estuarine, Coastal and Shelf Science* **86**: 71-82
- 65: Whanpetch, N., Nakaoka, M., Mukai, H., Suzuki, T., Nojima, S., Kawai, T. and Aryuthaka, C. (2010) Temporal changes in benthic communities of seagrass beds impacted by a tsunami in the Andaman Sea, Thailand. *Estuarine, Coastal and Shelf Science* **87**: 246-252
- 66: Okuda, T., Noda, T., Yamamoto, T., Hori, M. and Nakaoka, N. (2010) Contribution of environmental and spatial processes to rocky intertidal metacommunity structure. *Acta Oecologica* **36**: 413-422
- 67: Fukaya, K., Okuda, T., Nakaoka, M., Hori, M. and Noda, T. (2010) Seasonality in the strength and spatial scale of processes determining intertidal barnacle population growth. *Journal of Animal Ecology* **79**: 1270-1279
- 68: Okuda, T., Noda, T., Yamamoto, T., Hori, M. and Nakaoka, N. (2010) Relative contribution of environmental and spatial processes on determining community structure: A comparison among taxa in rocky intertidal assemblages. *Japanese Journal of Ecology* **60**: 227-239
- 69: Adachi, H., Yamano, H., Miyajima, T. and Nakaoka, M. (2010) A simple and robust procedure for coring unconsolidated sediment in shallow water. *Journal of Oceanography* **66**: 865-872
- 70: Yamada, K., Hori, M., Nakaoka, M. and Hamaguchi M. (2010) Temporal and spatial variations in functional-trait composition (functional diversity) of macrocrustacean communities in seagrass meadows. *Crustaceana Monographs* (Special edition: New Frontiers in Crustacean Biology) **15**: 325-339
- 71: Tanaka, N., Demise, T., Ishii, M., Shoji, Y. and Nakaoka, M. (2011) Genetic structure and gene flow of eelgrass *Zostera marina* populations in Tokyo Bay, Japan: implications for their restoration. *Marine Biology* **158**: 871-882
- 72: Yamakita, T., Wanatabe, K. and Nakaoka, M. (2011) Asynchronous local dynamics contributes to stability of a seagrass bed in Tokyo Bay. *Ecography* **34**: 519-528
- 73: Kubicek, A., Bessho, K., Nakaoka, M., Wahl, M. and Lenz M. (2011) Inducible defence and its modulation by environmental stress in the red alga *Chondrus yendoi* (Yamada and Mikami, 1965) from Honshu Island, Japan. *Journal of Experimental Marine Biology and Ecology* **397**: 203-213
- 74: Wahl, M., Link, H., Alexandridis, N., Thomason, J. C., Cifuentes, M., Costello, M. J., da Gama, B. A. P., Hillock, K., Hobday, A. J., Kaufmann, M. J., Keller, S., Kraufvelin, P., Kruger, I., Lauterbach, L., Antunes, B. L., Molis, M., Nakaoka, M., Nyström, J., bin Radzi, Z., Stockhausen, B., Thiel, M., Vance, T., Weseloh, A., Whittle, M., Wiesmann, L., Wunderer, L., Yamakita, T. and Lenz, M. (2011) Re-structuring of marine communities exposed to environmental change: a global study on the interactive effects of species and functional richness. *PLoS ONE* 6(5): e19514. doi:10.1371/journal.pone.0019514
- 75: Yamakita, T. and Nakaoka, M. (2011) Importance of considering grain and extent for the

analysis on spatial dynamics: perspectives from comparison between theory and empirical example on seagrass bed dynamics in Tokyo Bay. *Procedia Social and Behavioral Sciences* **21**: 177-183

- 76: Tanaka, N., Nakaoka, M., Omori, Y. and Aioi, K. (2012) Genetic diversity and gene flow of East Asian seagrass, *Zostera caespitosa* Miki (Zosteraceae), in northeast Japan. *Bulletin of the National Museum of Nature and Science. Series B* **38**: 29-36
- 77: Shimabukuro, H., Hori, M., Yoshimitsu, T., Tokunaga, N., Inokari, T., Sasaki, K., Nakaoka, M., Kawane, M., Yoshida, G. and Hamaguchi, M. (2012) Genetic differentiation of annual *Zostera marina* L. growing in Kagoshima Bay, Kagoshima, Japan based on an analysis using microsatellite markers. *Nippon Suisan Gakkaishi* **78**:204-211 (published March 2012)
- 78: Nakajima, Y., Matsuki, Y., Lian, C., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M. and Nadaoka, K. (2012) Development of novel microsatellite markers in a tropical seagrass, *Enhalus acoroides*. *Conservation Genetics Resources* **4**: 515-517. DOI 10.1007/s12686-012-9614-9 (published June 2012)
- 79: Fortes, M. D., Go, G. A., Bolisay, K., Nakaoka, M., Uy, W. H., Lopez, M. R., Leopardas, V., Leriorato, J., Allyn Pantallano, A., Paciencia Jr., F., Watai, M., Honda, K., and Edralin, M. (2012) Seagrass response to mariculture-induced physico-chemical gradients in Bolinao, northwestern Philippines. Proceedings of the 12th International Coral Reef Symposium, Cairns, Australia, 9-13 July 2012, ICRS2012_15B_3. (Published June 2012)
- 80: Kamauchi, H., Sato, S., Hayashi, D., Okabe, Y., Katsuyama, T., Fukushima, K., Yoshioka, A., Sato, T., Tokuchi, N. and Nakaoka, M. (2012) Consumer of salmon carcasses in early winter of Eastern Hokkaido. *Forest Research* **78**: 81-87
- 81: Kyoda, J., Hosokawa, S., Watanabe, K., Nakaoka, M., Shibamura, S., Montani, S. and Kuwae, T. (2012) Estimation of the distribution and coverage of seagrasses using field observation data and satellite imagery. *Journal of Hydraulic, Coastal and Environmental Engineering* **68**(2): I_1466-I_1470
- 82: Matsuki, Y., Nakajima, Y., Lian, C., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M. and Nadaoka, K. (2012) Development of microsatellite markers for *Thalassia hemprichii* (Hydrocharitaceae), a widely distributed seagrass in the Indo-Pacific. *Conservation Genetics Resources*. **4**: 1007–1010. DOI 10.1007/s12686-012-9694-6 (published December 2012)
- 83: Maeda, T., Kawai, T., Nakaoka, M. and Yotsukura, N. (2013) Effective DNA extraction method for fragment analysis using capillary sequencer of the kelp, *Saccharina*. *Journal of Applied Phycology*.**25**: 337-347. DOI 10.1007/s12686-013-9889-5 (Published February 2013)
- 84: Fukaya, K., Okuda, T., Hori, M., Yamamoto, T., Nakaoka, M. and Noda, T. (2013) Variable processes that determine population growth and invariant mean-variance relationship of intertidal barnacles. *Ecosphere* **4**:art48 DOI 10.1890/ES12-00272.1 (published April 2013)
- 85: Honda, K., Nakamura, Y., Nakaoka, M., Uy, W.H., Fortes, M.D. (2013) Habitat use by fishes in coral reefs, seagrass beds and mangrove habitats in the Philippines. *PLoS ONE* **8**: e65735 DOI 10.1371/journal.pone.0065735 (published August 2013)
- 86: Matsuki, Y., Takahashi, A., Nakajima, Y., Lian, C., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M. and Nadaoka, K. (2013) Development of microsatellite markers in a tropical seagrass *Syringodium isoetifolium* (Cymodoceaceae). *Conservation Genetic Resources*. DOI 10.1007/s12686-013-9889-5 (published September 2013)
- 87: Nakajima, Y., Yasuda, N., Matsuki, Y., Arriessgado, D.M., Lian, C., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M., Taquet, C., Suharsono, Nagai, S. and Nadaoka, K. (2013) Development of 10 novel polymorphic microsatellite markers for the Indo-Pacific horned starfish, *Protoreaster nodosus*. *Marine Genomics* **11**: 27-29 DOI 10.1016/j.margen.2013.05.002 (published September 2013)
- 88: Arriessgado, D.M., Nakajima, Y., Matsuki, Y., Lian, C., Nagai, S., Yasuike, N., Nakamura, Y., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M. and Nadaoka, K. (2014) Development of novel microsatellite markers for *Cymodocea rotundata* Ehrenberg (Cymodoceaceae), a pioneer species widely distributed in the Indo-Pacific. *Conservation Genetic Resources*.**6**: 135-138 DOI 10.1007/s12686-013-0025-3 (published March 2014)
- 89: Sato, M., Honda, K., Bolisay, K.O., Nakamura, Y., Fortes, M.D. and Nakaoka, M. (2014)

- Factors affecting the local abundance of two anemonefishes (*Amphiprion frenatus* and *A. perideraion*) around a semi-closed bay in Puerto Galera, the Philippines. *Hydrobiologia* **733**: 63-69 (published July 2014)
- 90: Rashidul Alam, A.K.M, Hagino, T., Fukaya, K., Okuda, T., Nakaoka, M. and Noda T. (2014) Early phase of the invasion of *Balanus glandula* along the coast of eastern Hokkaido: changes in abundance, distribution and recruitment. *Biological Invasions* **16**: 1699-1708 (published August 2014)
- 91: Leopardas, V.E., Uy, W., Nakaoka, M. (2014) Benthic macrofaunal assemblages in multispecific seagrass meadows of the southern Philippines: Variation among vegetation dominated by different seagrass species. *Journal of Experimental Marine Biology and Ecology* **457**: 71–80 (published August 2014)
- 92: Yamada, K., Tanaka, Y., Era, T. and Nakaoka, M. (2014) Environmental and spatial controls of macroinvertebrate assemblages in seagrass ecosystems. *Global Ecology and Conservation* **2**: 47-62 (Available online 21 August 2014)
- 93: Sato, M., Kurokochi, H., Tan, E., Asakawa, S., Honda, K., Bolisay, K.O., Nakamura, Y., Lian, C., Fortes, M.D. and Nakaoka, M. (2014) Fifteen novel microsatellite markers for two Amphiprion species (*Amphiprion frenatus* and *Amphiprion perideraion*) and cross-species amplification. *Conservation Genetics Resources* **6**: 685–688 (published September 2014)
- 94: Tanaka, Y., Go, G.A., Watanabe, A., Miyajima, T., Nakaoka, M., Uy, W.H., Nadaoka, K., Watanabe, S., Fortes, M.D. (2014) 17-year change in species composition of mixed seagrass beds around Santiago Island, Bolinao, the northwestern Philippines. *Marine Pollution Bulletin* **88**: 81-85 (published November 2014) DOI: 10.1016/j.marpolbul.2014.09.024
- 95: Fukaya, K., Okuda, T., Nakaoka, M. and Noda, T. (2014) Effects of spatial structure of population size on the population dynamics of barnacles across their elevational range. *Journal of Animal Ecology* **83**: 1334–1343 (published November 2014) DOI: 10.1111/1365-2656.12234
- 96: Nakajima, Y., Matsuki, Y., Lian, C., Fortes, M.D., Uy, W.H., Campos, W.L., Nakaoka, M. and Nadaoka K. (2014) The Kuroshio Current influences genetic diversity and population genetic structure of a tropical seagrass, *Enhalus acoroides*. *Molecular Ecology* **23**: 6029–6044 (published December 2014), DOI: 10.1111/mec.12996
- 97: Yamakita, T., Yamamoto, H., Nakaoka, M., Yamano, H., Fujikura, K., Hidaka, K., Hirota, Y., Ichikawa, T., Kakehi, S., Kameda, T., Kitajima, S., Kogure, K., Komatsu, T., Kumagai, N. H., Miyamoto, H., Miyashita, K., Morimoto, H., Nakajima, R., Nishida, S., Nishiuchi, K., Sakamoto, S., Sano, M., Sudo, K., Sugisaki, H., Tadokoro, K., Tanaka, K., Jintsu-Uchifune, Y., Watanabe, K., Watanabe, H., Yara, Y., Yotsukura, N. and Shirayama, Y. (2015) Identification of important marine areas around the Japanese Archipelago: Establishment of a protocol for evaluating a broad area using ecologically and biologically significant areas selection criteria. *Marine Policy* **51**: 136-147 (published January 2015) DOI: 10.1016/j.marpol.2014.07.009
- 98: Watai, M., Nakamura, Y., Honda, K., Bolisay, K.O., Miyajima, T., Nakaoka, M. and Fortes, M.D. (2015) Diet, growth, and abundance of two seagrass bed fishes along a pollution gradient caused by milkfish farming in Bolinao, northwestern Philippines. *Fisheries Science* **81**: 43-51 (published January 2015) DOI: 10.1007/s12562-014-0824-9
- 99: Hosokawa, S., Nakaoka, M., Miyoshi, E. and Kuwae, T. (2015) Seed dispersal in the seagrass *Zostera marina* L. is mostly within the parent bed in a protected bay. *Marine Ecology Progress Series* **523**: 41–56 (published March 16, 2015) DOI:10.3354/meps11146
- 100: Miyajima, T., Hori, M., Hamaguchi, M., Shimabukuro, H., Adachi, H., Yamano, H. and Nakaoka, M. (2015) Geographic variability in organic carbon stock and accumulation rate in sediments of East and Southeast Asian seagrass meadows. *Global Biogeochemical Cycles* **29**: 397-415 (published online: 7 April 2015) DOI: 10.1002/2014GB004979
- 101: Duffy, J.E., Reynolds, P.L., Boström, C., Coyer J.A., Cusson, M., Donadi, S., Douglass, J.A., Eklöf, J.S., Engelen, A.H., Eriksson, B.K., Fredriksen, S., Gamfeldt, L., Gustafsson, C., Hoarau, G., Hori, M., Hovel, K., Iken, K., Lefcheck, J.S., Moksnes, P-O., Nakaoka, M., O'Connor, M.I., Olsen, J.L., Richardson, J.P., Ruesink, J.L., Sotka, E.E., Thormar, J., Whalen,

- M.A. and Stachowicz, J.J. (2015) Biodiversity mediates top-down control in eelgrass ecosystems : a global comparative-experimental approach. *Ecology Letters* **18**: 696–705 (published in 17 May 2015) DOI: 10.1111/ele.12448
- 102: Honda, K., Uy, W.H., Baslot, D.I., Pantallano, A.D.S., Nakamura, Y., and Nakaoka, M. (2016) Diel habitat-use patterns of fishery target fishes in a marine protected area in the Philippines. *Aquatic Biology* Vol. **24**: 163–174 (published in January 27, 2016) doi: 10.3354/ab00646
- 103: Sahara, R., Fukaya, K., Okuda, T., Hori, M., Yamamoto, T., Nakaoka, M. and Noda, T. (2016) Larval dispersal dampens population fluctuation and shapes the interspecific spatial distribution patterns of rocky intertidal gastropods. *Ecography* **39**: 487-495 (published online on July 14, 2015) DOI: 10.1111/ecog.01354
- 104: van Katwijk, M.M., Thorhaug, A., Marbà, N., Orth, R.J., Duarte, C.M., Kendrick, G.A., Althuisen, I.H.J., Balestri, E., Bernard, G., Cambridge, M.J., Cunha, A., Durance, C., Giesen, W., Han, Q., Hosokawa, S., Kiswara, W., Komatsu, T., Lardicci, C., Lee, K-S., Meinesz, A., Nakaoka, M., O'Brien, K., Paling, E.I., Pickerell, C., Ransijn, A.M.R. and Verduin, J.J. (2016) Global analysis of seagrass restoration: the importance of large-scale planting. *Journal of Applied Ecology* **53**: 567-578 DOI: 10.1111/1365-2664.12562
- 105: Yotsukura, N., Maeda, T., Abe, T., Nakaoka, M. and Kawai, T. (2016) Genetic differences among varieties of *Saccharina japonica* in northern Japan as determined by AFLP and SSR analyses. *Journal of Applied Phycology* **28**: 3043–3055 (accepted on January 21, 2016; published online on February 13, 2016) DOI: 10.1007/s10811-016-0807-6
- 106: Leopardas, V. Honda, K., Go, G.A., Bolisay, K., Pantallano, A.D., Uy, W., Fortes, M.D. and Nakaoka, M. (2016) Variation in macrofaunal community of sea grass beds along a pollution gradient in Bolinao, northwestern Philippines. *Marine Pollution Bulletin* **105**: 310–318 (published online on February 15, 2016) DOI: 10.1016/j.marpolbul.2016.02.004
- 107: Maass, M., Balvanera, P., Baudry, J., Bourgeron, P., Dick, J., Equihua, M., Forsius, M., Halada, L., Krauze, K., Nakaoka, M., Orenstein, D.E., Parr, T., Redman, C.L., Rozzi, R., Santos-Reis, M., Swemmer, T. and Vădineanu, A. (2016) Changes in biodiversity and trade-offs among ecosystem services, stakeholders and components of well-being: the contribution of the International Long-Term Ecological Research network (ILTER) to Programme on Ecosystem Change and Society (PECS). *Ecology and Society* **21**(3): 31 (accepted on May 30, 2016; published online on September 12, 2016) DOI: 10.5751/ES-08587-210331
- 108: Fukaya, K., Royle, J.A., Okuda, T., Nakaoka, M., Noda, T. (2017) A multistate dynamic site occupancy model for spatially aggregated sessile communities. *Methods in Ecology and Evolution* **8**: 757–767 (accepted on October 10, 2016; Published online on November 28, 2016) DOI: 10.1111/2041-210X.12690
- 109: Momota, K. and Nakaoka, M. (2017) Influence of different types of sessile epibionts on the community structure of mobile invertebrates in an eelgrass bed. *PeerJ* **5**: e2952; DOI: 10.7717/peerj.2952 (accepted on January 2, 2017, published online on January 31, 2017)
- 110: Fukuda, A., Kamimura, Y., Hori, M., Nakaoka, M., Noda, T., Yamashita, Y., Otake, T. and Shoji, J. (2017) Offshore currents explain the discontinuity of fish community in seagrass bed along the Japanese archipelago. *Fisheries Oceanography* **26**: 65-68 DOI: doi:10.1111/fog.12186 (accepted on August 26, 2016; Published online on April 2017)
- 111: Wang, S., Wang, G., Weinberger, F., Bian, D., Nakaoka, M. and Lenz, M. (2017) Anti-epiphyte defences in the red seaweed *Gracilaria vermiculophylla*: non-native algae are better defended than their native conspecifics. *Journal of Ecology* **105**: 445-457 (accepted on October 20, 2016, published online on November 28, 2016) DOI: 10.1111/1365-2745.12694
- 112: Shao, H., Minami, K., Shirakawa, H., Maeda, T., Ohmura, T., Fujikawa, Y., Yotsukura, N., Nakaoka, M., and Miyashita, K. (2017) Verification of echosounder measurements of thickness and spatial distribution of kelp forests. *Journal of Marine Science and Technology* **25**: 343-351 (accepted on December 29, 2016, published in May 2017) DOI: 10.6119/JMST-016-1229-1
- 113: Honda, K., Uy, W.H., Baslot, D.I., Pantallano, A.D.S., Sato, M., Nakamura, Y. and Nakaoka, M. (2017) Importance of outer reef slopes for commercially important fishes: implications for designing a marine protected area in the Philippines. *Fisheries Science* **83**: 523-535

- (published online on April 4, 2017, accepted on March 17, 2017) DOI: 10.1007/s12562-017-1082-4
- 114: Sharma, S., Nadaoka K., Nakaoka, M., Uy, W.H., MacKenzie, R.A., Friess, D.A., and Fortes, M.D. (2017) Growth performance and structure of a mangrove afforestation project on a former seagrass bed, Mindanao Island, Philippines. *Hydrobiologia* **803**: 359-371 (published online on June 5, 2017, accepted on May 28, 2017) DOI 10.1007/s10750-017-3252-x
- 115: Sato, M., Honda, K., Uy, W.H., Baslot, D.I., Genovia, T.G., Nakamura, Y., Bernardo, L.P.C., Kurokochi, H., Pantallano, A.D.S., Lian, C., Nadaoka, K. and Nakaoka, M. (2017) Marine protected area restricts demographic connectivity: dissimilarity in a marine environment can function as a biological barrier. *Ecology and Evolution* **7**: 7859-7871 (accepted on July 15, 2017, published online on August 30, 2017) DOI: 10.1002/ece3.3318
- 116: Wang, S., Weinberger, F., Xiao, L., Nakaoka, M., Wang, G., Krueger-Hadfield, S.A., Sotka, E.E., Bian, D. and Lenz, M. (2017) In-situ-common-garden-assays demonstrate increased defense against natural fouling in non-native populations of the red seaweed *Gracilaria vermiculophylla*. *Marine Biology* **164**: 193 (accepted on August 24, 2017, published online on September 5, 2017) DOI: 10.1007/s00227-017-3226-6
- 117: Ruesink, J., Stachowicz, J.J., Reynolds, P., Boström, C., Cusson, M., Douglass, J., Eklöf, J., Engelen, A. Hori, M., Hovel, K., Iken, K., Moksnes, P., Nakaoka, M., O'Connor, M., Olsen, J., Sotka, E., Whalen, M. and Duffy, E. (2017) Form-function relationships in a marine foundation species depend on scale: a shoot to global perspective from a distributed ecological experiment. *Oikos* (accepted on August 21, 2017) DOI: 10.1111/oik.04270
- 118: Reynolds, P.L., Stachowicz, J.J., Hovel, K., Boström, C., Boyer, K., Cusson, M., Eklöf, J.S., Engel, F.G., Engelen, A. H., Klemens, B.K., Fodrie, F.J., Griffin, J.N., Hereu, C., Hori, M., Hanley, T., Ivanov, M., Jorgensen, P. Kruschel, C., Lee, K-S., McGlathery, K., Moksnes, P.O., Nakaoka, M., Nash, F.T., O'Connor, M.I., O'Connor, N., Orth, R.J., Rossi, F., Ruesink, J., Sotka, E., Unsworth, R.K.F., Whalen, M.A., Duffy, J.E. (in press) Latitude, temperature and habitat complexity predict predation pressure in eelgrass beds across the Northern Hemisphere. *Ecology* (accepted on August 30, 2017) DOI: 10.1002/ecy.2064
- 119: Nordlund, L.M., Jackson, E.L., Nakaoka M., Samper-Villarreal, J., Beca-Carretero, P. and Creed, J.C. (in press) Seagrass ecosystem services – what's next? *Marine Pollution Bulletin* (accepted on September 7, 2017) DOI: 10.1016/j.marpolbul.2017.09.014
- 120: Leopardas, V. E., Hori, M., Mukai, H., Tanaka, Y., Yokoi, K., Aoki, M., Sato, N., Whanpetch, N. and Nakaoka, M. (in press) Broad scale variation in seagrass benthic macrofaunal assemblages along the coast of Japan. *Ecological Research* (accepted on October 7, 2017) DOI: 10.1007/s11284-017-1517-5
- 121: Nakaoka, M., Sudo, K., Namba, M., Shibata, H., Nakamura, F., Ishikawa, S., Makino, M., Yamano, H., Matsuzaki, S.S., Yamakita, T., Yu, X., Hou, X., Li, X., Brodie, J., Kanemoto, K., Moran, D. and Verones, F. (in press) TSUNAGARI: A new interdisciplinary and transdisciplinary study toward conservation and sustainable use of biodiversity and ecosystem services. *Ecological Research* (accepted on October 25, 2017) DOI: 10.1007/s11284-017-1534-4
- 122: Fang, X., Hou, X., Li, X., Hou, W., Nakaoka, M. and Yu, X (in press) Ecological connectivity between land and sea: a review. *Ecological Research* (accepted on December 8, 2017) DOI: 10.1007/s11284-017-1549-x
- 123: Sotka, E.E., Baumgardner, A.W., Bippus, P., Destombe, C., Duermit, E., Endo, H., Flanagan, B.A., Kamiya, M., Lees, L., Murren, C.J., Nakaoka, M., Shainker, S., Strand, A.E., Terada, R., Valero, M., Weinberger, F. and Krueger-Hadfield, S.A. (in press) Combining niche-shift and population genetic analyses predicts rapid phenotypic evolution during invasion. *Evolutionary Applications* (accepted on December 26, 2017)

Books

- 1: Aioi, K. and Nakaoka, M. (2003) Seagrasses of Japan. In, Green, E.P. and Short, F.T., editors, *World Atlas of Seagrasses*. University of California Press, Berkeley, USA, pp. 185-192
- 2: Yamamoto, T., Komatsu T., Kawai H., Nakaoka, M., Marine Life Research Group of Takeno and

- Ohwada, K. (2003) Impact and recovery of an intertidal ecosystem from the *Nakhodka* heavy-oil spill. In, Tazaki, K., editor, Heavy Oil Spilled from Russian Tanker “*Nakhodka* “ in 1997: Toward Eco-responsibility, Earth Sense. 21st Century COE Kanazawa University, Kanazawa, Japan, pp. 162-173
- 3: Nakaoka, M. (2003) Population Ecology and Life History Study. In, Wada, K., editor, *Ecology of Marine Benthos*, Tokai University Press, Kanagawa, pp. 33-115
 - 4: Nakaoka, M. (2005) Biodiversity of seagrass beds. In, Miyazaki, N., editor, Sea of Sanriku, Scientist Co. Ltd., Tokyo, pp. 66-83
 - 5: Komatsu, T., Nakaoka, M., Kawai, H., Yamamoto, T., Marine Life Research Group of Takeno and Ohwada, K. (2005) Impact assessment of the heavy-oil pollution on the intertidal ecosystems after the ‘*Nakhodka*’ oil spill by GIS. In, Ohwada, K., Koyama, J. and Hiroishi S., editors, Effect of Oil Spill Accidents onto Marine Environments with Special Attention to *Nakhodka* Case, Koseisha-Koseikaku, Tokyo, pp. 120-134
 - 6: Nakaoka, M., Kouchi, N., Yoshida, M., Ohno, M. and Kobayashi, A. (2008) Jangusa Watch Handbook., Nature Conservation Society of Japan, Tokyo, 36 pp
 - 7: Nakaoka, M. (2008) Changes in coastal ecosystems in relation to global climate change, with special reference to biological community dynamics. In, Ohgushi, T., Kondoh, M. and Nakaoka, M., editors, Community Ecology Series 4: Linking Ecosystem with Community. Kyoto University Press, Kyoto, pp. 179-204
 - 8: Nakaoka, M., Kondoh, M. and Ohgushi T. (2008) Linking biological community and ecosystem. In, Ohgushi, T., Kondoh, M. and Nakaoka, M., editors, Community Ecology Series 4: Linking Ecosystem with Community. Kyoto University Press, Kyoto, pp. 205-219
 - 9: Nakaoka, M. (2010) Biodiversity and function of seagrass bed. In, Ecological Society of Japan, editor, Ecology Lecture Series 3: Why shall we protect biodiversity on earth? Bun-ichi Sogo Press, Tokyo, pp. 6-17
 - 10: Ecological Society of Japan, Nakaoka, M. editors (2011) Ecology Lecture Series 4: Challenges of ecology toward global environmental problems. Bun-ichi Sogo Press, Tokyo
 - 11: Nakaoka, M. and Matsuda, H. (2011) Elucidating the link among global climate change, ecosystem service and human activity. In, Shoji, J., Hori, M. and Yamashita, Y., editors, Ecosystem Services of Coastal Waters: Benefits of the Sea and its Sustainable Use. Kouseisha Kouseikaku, Tokyo, pp. 129-143
 - 12: Sato, H., Shimada, M., Takemon, Y., Nakaoka, M. and Muraoka, H. (2012) Conservation of ecosystems and earth environment. In, Ecological Society of Japan, editor, Introduction to Ecology 2nd edition, Tokyo Kagaku Dojin, Tokyo, pp. 227-263
 - 13: Nakaoka, M. (2012) Macrophyte beds. In, Shirayama, Y., Sakurai, Y., Furuya, K., Nakahara, H., Matsuda, H. and Kagami, Y., editors, Marine Conservation Ecology. Kodansha, Tokyo, pp. 66-68
 - 14: Ohte, N., Nakaoka, M. and Shibata H. (2012) ILTER and JaLTER: their missions and linkage to database development in the Asia-Pacific region. In: Nakano S., Yahara T., Nakashizuka T. (eds) The Biodiversity Observation Network in the Asia-Pacific Region. Ecological Research Monographs. Springer, Tokyo, pp. 205-215
 - 15: Nakaoka, M., Lee, K-S., Huang, X., Almonte, T., Sidik Bujang, J., Kiswara, W., Ambo Rappe, R., Maryam Yaakub, S., Prabhakaran, M.P., Abu Hena M.K., Hori, M., Zhang, P., Prathep, A. and Fortes, M. D. (2014) Regional comparison of the ecosystem services from seagrass beds in Asia. In: (S. Nakano et al. eds.) Asia-Pacific Biodiversity Observation Network: Integrative Observations and Assessments. Ecological Research Monographs, Springer, Tokyo, pp. 367-391 (published May 6, 2014)
 - 16: Nakaoka, M. (2016) Diversity of marine organisms. In, Ecological Society of Japan, editor, Current Ecology Series 10: Marine Ecology, Kyoritsu Shuppan, Tokyo, pp. 7-20
 - 17: Kawamura, T. and Nakaoka, M. (2016) Sea-bottom ecosystems. In, Ecological Society of Japan, editor, Current Ecology Series 10: Marine Ecology, Kyoritsu Shuppan, Tokyo, pp. 68-121

Others

- 1: Nakaoka, M. (1989) The growth and reproductive cycles of *Yoldia* sp. [aff. *Y. notabilis*

- Yokoyama] (Mollusca; Bivalvia) in Otsuchi Bay. *Otsuchi Marine Research Center* **15**: 21-27
- 2: Nakaoka, M. (1993) Influence of variation in larval settlement processes on the population dynamics of marine benthic animals: analyses using projection matrix models. *Gekkan Kaiyo* **25**: 264-268
 - 3: Nakaoka, M. and Matsumasa, M. (1997) Mechanisms of creating community diversity: cases on benthic communities. *Journal of Plankton Society of Japan* **44**: 37-38
 - 4: Komatsu, T., Nakaoka, M. and Kawai, H. (1998) Long-term monitoring of coastal ecosystems polluted by heavy-oil spill using Geographical Information System (GIS). *Gekkan Kaiyo* **30**: 648-653
 - 5: Komatsu, T., Nakaoka, M., Kawai, H., Kamiya, Y. and T. Yamamoto (2000) Long-term effects of oil spill on intertidal ecology. *Yudaku Kikin Dayori* **67**: 1-11
 - 6: Nakaoka, M. (2000) Small invertebrate animals using flowers and seeds of seagrasses. *Umiushi-Tushin* **27**: 9-11
 - 7: Nakaoka, M., Supanwanid C (2000) Quantitative estimation of the distribution and biomass of seagrass at Haad Chao Mai National Park, Trang Province, Thailand. *Kasetsart University Fishery Research Bulletin* **22**: 10-22
 - 8: Watanabe, M., Nakaoka, M. (2000) Environmental and biological factors affecting distribution and productivity of seagrasses. *Aquabiology* **22**: 533-541
 - 9: Toyohara, T., Kouchi, N., Nakaoka, M. (2000) Ecology of epiphytic animals in seagrass beds. *Aquabiology* **22**: 557-565
 - 10: Nakaoka, M. (2000) Development of local and global networks for research and conservation of seagrasses. *Aquabiology* **22**: 566-569
 - 11: Kawai, H., Kamiya, Y., Komatsu, T., Nakaoka, M., Yamamoto, T., Marine Life Research Group of Takeno (2001) Impacts by heavy-oil spill from the Russian tanker "Nakhodka" on intertidal ecosystems: from an impact assessment of the oil spill to a long-term monitoring program. *Iden* **55** (4): 92-97
 - 12: Nakaoka, M. (2002) General characteristics of exotic species revealed by the studies of marine bivalve *Musculista senhousia*: Interpretation and comments on the lecture by Dr Jef Crooks. *Annual report of Kanto Regional Branch of Japanese Society of Ecology* **50**: 24-27
 - 13: Nakaoka, M. (2002) Effects of material input from terrestrial ecosystems on the dynamics of tropical seagrass community: comparisons of seagrass beds with different impact of rivers. *Gekkan Kaiyo* **34**: 424-429
 - 14: Nakaoka, M., Noda, T. (2004) Special feature: multiple spatial scale approaches in population and community ecology. *Population Ecology* **46**: 103-104
 - 15: Nakaoka, M. (2005) On food-web flexibility: beyond the gap between theoretical and empirical studies. *Japanese Journal of Ecology* **55**: 265-268
 - 16: Nakaoka, M., Kouchi, N., Yoshida, M., Kobayashi, A. (2007) Jangusa Watch: Census of seagrass bed based on citizen-based monitoring. In, Report of Seagrass Monitoring Survey at Eastern Shore of Northern Okinawa. Nature Conservation Society of Japan, Tokyo, pp.3-31
 - 17: Yoshida, M., Kouchi, N., Nakaoka, M. (2007) Comparisons of remote sensing survey of seagrass bed by an aerial blimp and citizen-based monitoring. In, Report of Seagrass Monitoring Survey at Eastern Shore of Northern Okinawa. Nature Conservation Society of Japan, Tokyo, pp.33-41
 - 18: Nakaoka, M., Ishibashi, C., Kouchi, N., Yoshida, M. (2007) Status of seagrass bed around Henoko area. In, Report of Seagrass Monitoring Survey at Eastern Shore of Northern Okinawa. Nature Conservation Society of Japan, Tokyo, pp.43-48
 - 19: Nakaoka, M. (2007) Status of seagrass in the planning area of land reclamation for the construction project of substitute airway for Futenma Airfield (short notes). In, Report of Seagrass Monitoring Survey at Eastern Shore of Northern Okinawa. Nature Conservation Society of Japan, Tokyo, pp.61-63
 - 20: Nakaoka, M. (2008) Current status and prospect of long-term monitoring of coastal ecosystems. In, *Current Status and Prospect of Long-term Ecosystem Monitoring: Influence of Global Warming and Responses of Ecosystems*. *Earth Environmental Center Report*, National Institute

- of Environmental Science, Tsukuba, pp. 59-62
- 21: Noda, T., Nakaoka, M., Takada, T. (2009) Special feature: Spatial connectivity and scaling for populations and communities. *Population Ecology* **51**: 3-4
 - 22: Nakaoka, M. (2009) Ecosystem and biodiversity monitoring in coastal area. *Iden* **63**(6): 80-86
 - 23: Nakaoka, M. (2010) Worldwide decline of seagrass bed revealed by global database analysis. *Iden* **64**(3): 85-86
 - 24: Hosokawa, S., Nakaoka, M., Miyoshi, E., Kuwae, T. (2010) Expansion of eelgrass distribution and the role of seed dispersal. *Report of Port and Airport Research Institute* **43**: 31-49
 - 25: Nakaoka, M., Watanabe, K. (2011) Biodiversity and ecosystem monitoring of seagrass beds. *Aquabiology* **33**(4): 315-322
 - 26: Yoshida, M., Abe, M., Ohno, M., Shikatani, M., Kouchi, N. and Nakaoka, M. (2013) Temporal change in seagrass bed at Henoko, Nago, Okinawa Island. Report of the Nature Conservation Society of Japan No. 100: 10-year report of "Jangusa Watch", Seagrass bed monitoring at Henoko, Okinawa Island. pp. 6-20 (published September 2013)
 - 27: Nakaoka, M., Suzuki, T., Dasai, A., Sakanishi, Y., Kurashima, A., Aoki, M. and Tanaka, J. (2013) Effects of the 2011 off the Pacific coast of Tohoku earthquake on coastal ecosystems. 2008-2012 Summary Report of Monitoring Sites 1000 Coastal Area Survey (Rocky Intertidal, Tidal Flats, Seagrass Beds, Algal Beds). Biodiversity Center of Japan, Nature Conservation Bureau, Ministry of the Environment, Japan. pp. 74-80
 - 28: Nakaoka, M., Hori, M., Tanaka, Y. and Mukai, H. (2013) Seagrass bed ecosystem. 2008-2012 Summary Report of Monitoring Sites 1000 Coastal Area Survey (Rocky Intertidal, Tidal Flats, Seagrass Beds, Algal Beds). Biodiversity Center of Japan, Nature Conservation Bureau, Ministry of the Environment, Japan. pp. 31-47
 - 29: Nakaoka, M., Isada, T., Honda, K. and Kitamura, T. (2014) Akkeshi Marine Station, Hokkaido University (From field research sites (26)). *Japanese Journal of Ecology* **64**: 87-91 (published March 30, 2014)
 - 30: Enoki, T., Nakashizuka, T., Nakano, S., Miki, T., Lin, Y. P., Nakaoka, M., Mizumachi, E. and Shibata, H. (2014) Progress in the 21st century: A Roadmap for the Ecological Society of Japan. *Ecological Research* **29**: 357-368 (published May 2014)
 - 31: Nakaoka, M., Tamaki, H., Muraoka, D., Tokuoka, M., Komatsu, T. and Tanaka, N. (2017) Temporal change in seagrass beds of Sanriku Coast before and after the Great East Japan Earthquake. *Nippon Suisan Gakkaishi* **83**: 659-663 (published August 2017)

Master & Ph.D theses

- 1: Nakaoka, M. (1989) Ecological study on the seasonality in the reproduction of *Limopsis tajimae* Sowerby (Bivalvia, Limopsidae) from the bathyal zone of Suruga Bay, Central Japan. Master's dissertation, University of Tokyo. 48pp.
- 2: Nakaoka, M. (1993) Population dynamics and life history traits of *Yoldia notabilis* (Bivalvia: Protobranchia). Doctoral dissertation, University of Tokyo. 182pp.